

APPLICANTS:
U.S.S.N.:

Gerlach et al.
09/898,586

Gerlach

(c) determining the presence or amount of said probe bound to said nucleic acid molecule,
thereby determining the presence or amount of the nucleic acid molecule in said sample.

- C2*
80. (New) The method of claim 19, wherein said nucleic acid molecule further comprises nucleotides 35-54 of SEQ ID NO:23, or a variant thereof, wherein said variant comprises a nucleic acid wherein one nucleotide is changed from nucleotides 35-45 of SEQ ID NO:23.
81. (New) The method of claim 19, wherein said nucleic acid molecule further comprises nucleotides 25-54 of SEQ ID NO:23, or a variant thereof, wherein said variant comprises a nucleic acid wherein one nucleotide is changed from nucleotides 25-54 of SEQ ID NO:23.
82. (New) The method of claim 19, wherein said nucleic acid molecule further comprises nucleotides 15-54 of SEQ ID NO:23, or a variant thereof, wherein said variant comprises a nucleic acid wherein one nucleotide is changed from nucleotides 15-54 of SEQ ID NO:23.
83. (New) The method of claim 19, wherein said nucleic acid molecule further comprises nucleotides 1-54 of SEQ ID NO:23, or a variant thereof, wherein said variant comprises a nucleic acid wherein one nucleotide is changed from nucleotides 1-54 of SEQ ID NO:23.
-